



SLC11A2 gene

solute carrier family 11 member 2

Normal Function

The *SLC11A2* gene provides instructions for making a protein called divalent metal transporter 1 (DMT1). Four different versions (isoforms) of the DMT1 protein are produced from the *SLC11A2* gene. Each isoform is specific to one or more tissues, but some form of the DMT1 protein is found in all tissues. The primary role of the DMT1 protein is to transport positively charged iron atoms (ions) within cells; however, the protein can transport some other metal ions as well.

In a section of the small intestine called the duodenum, the DMT1 protein is located within finger-like projections called microvilli. These projections absorb nutrients from food as it passes through the intestine and then release them into the bloodstream. In all other cells, including immature red blood cells called erythroblasts, DMT1 is located in the membrane of endosomes, which are specialized compartments that are formed at the cell surface to carry proteins and other molecules to their destinations within the cell. DMT1 transports iron from the endosomes to the cytoplasm so it can be used by the cell.

Health Conditions Related to Genetic Changes

hypochromic microcytic anemia with iron overload

At least seven mutations in the *SLC11A2* gene have been found to cause hypochromic microcytic anemia with iron overload. This condition is characterized by a shortage of red blood cells (anemia) that is apparent at birth. The red blood cells that are produced are abnormally small (microcytic) and pale (hypochromic). There is also progressive accumulation of iron in the liver.


Most *SLC11A2* gene mutations that cause this condition change single protein building blocks (amino acids) in the DMT1 protein. These mutations lead to reduced production of the DMT1 protein, decreased protein function, or impaired ability of the protein to get to the correct location in cells. In erythroblasts, a shortage of DMT1 protein diminishes the amount of iron transported within cells, even though there is an abundance of iron in the blood. As a result, the development of healthy red blood cells is impaired, leading to a shortage of these cells. In the duodenum, a shortage of DMT1 protein decreases iron absorption. To compensate, cells increase production of functional DMT1 protein, which increases iron absorption. Because the red blood cells cannot use the iron that is absorbed, it accumulates in the liver, eventually impairing liver function. The lack of iron in red blood cells and the accumulation of

iron in the liver lead to the signs and symptoms of hypochromic microcytic anemia with iron overload.

Chromosomal Location

Cytogenetic Location: 12q13.12, which is the long (q) arm of chromosome 12 at position 13.12

Molecular Location: base pairs 50,952,263 to 51,028,335 on chromosome 12 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- DCT1
- divalent cation transporter 1
- DMT-1
- DMT1
- natural resistance-associated macrophage protein 2
- NRAMP 2
- NRAMP2
- solute carrier family 11 (proton-coupled divalent metal ion transporter), member 2
- solute carrier family 11 (proton-coupled divalent metal ion transporters), member 2

Additional Information & Resources

Educational Resources

- Biochemistry (fifth edition, 2002): Oxygen Binding Induces Substantial Structural Changes at the Iron Sites in Hemoglobin
<https://www.ncbi.nlm.nih.gov/books/NBK22596/#A1348>
- Madame Curie Bioscience Database: A Glimpse into the Mechanism of Metal-Ion Uptake
<https://www.ncbi.nlm.nih.gov/books/NBK6091/#A23429>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28SLC11A2%5BTIAB%5D%29+OR+%28%28divalent+metal+transporter+1%5BTIAB%5D%29+OR+%28DMT1%5BTIAB%5D%29+OR+%28NRAMP2%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>

OMIM

- SOLUTE CARRIER FAMILY 11 (PROTON-COUPLED DIVALENT METAL ION TRANSPORTER), MEMBER 2
<http://omim.org/entry/600523>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_SLC11A2.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=SLC11A2%5Bgene%5D>
- HGNC Gene Family: Solute carriers
<http://www.genenames.org/cgi-bin/genefamilies/set/752>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=10908
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/4891>
- UniProt
<http://www.uniprot.org/uniprot/P49281>

Sources for This Summary

- Bardou-Jacquet E, Island ML, Jouanolle AM, D tivaud L, Fatih N, Ropert M, Brissot E, Mosser A, Maisonneuve H, Brissot P, Lor al O. A novel N491S mutation in the human SLC11A2 gene impairs protein trafficking and in association with the G212V mutation leads to microcytic anemia and liver iron overload. *Blood Cells Mol Dis.* 2011 Dec 15;47(4):243-8. doi: 10.1016/j.bcmd.2011.07.004. Epub 2011 Aug 26.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21871825>
- Beaumont C, Delaunay J, Hetet G, Grandchamp B, de Montalembert M, Tchernia G. Two new human DMT1 gene mutations in a patient with microcytic anemia, low ferritinemia, and liver iron overload. *Blood.* 2006 May 15;107(10):4168-70. Epub 2006 Jan 26.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16439678>

- Iolascon A, De Falco L. Mutations in the gene encoding DMT1: clinical presentation and treatment. *Semin Hematol*. 2009 Oct;46(4):358-70. doi: 10.1053/j.seminhematol.2009.06.005. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/19786204>
- Lam-Yuk-Tseung S, Camaschella C, Iolascon A, Gros P. A novel R416C mutation in human DMT1 (SLC11A2) displays pleiotropic effects on function and causes microcytic anemia and hepatic iron overload. *Blood Cells Mol Dis*. 2006 May-Jun;36(3):347-54. Epub 2006 Apr 3.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16584902>
- Mims MP, Prchal JT. Divalent metal transporter 1. *Hematology*. 2005 Aug;10(4):339-45. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16085548>
- Priwitzerova M, Nie G, Sheftel AD, Pospisilova D, Divoky V, Ponka P. Functional consequences of the human DMT1 (SLC11A2) mutation on protein expression and iron uptake. *Blood*. 2005 Dec 1; 106(12):3985-7. Epub 2005 Aug 9.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16091455>
- OMIM: SOLUTE CARRIER FAMILY 11 (PROTON-COUPLED DIVALENT METAL ION TRANSPORTER), MEMBER 2
<http://omim.org/entry/600523>
- Shawki A, Knight PB, Maliken BD, Niespodzany EJ, Mackenzie B. H(+)-coupled divalent metal-ion transporter-1: functional properties, physiological roles and therapeutics. *Curr Top Membr*. 2012;70: 169-214. doi: 10.1016/B978-0-12-394316-3.00005-3. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/23177986>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/SLC11A2>

Reviewed: November 2014
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services